

AN EXTENSION OF HARMONIC FUNCTIONS ALONG FIXED DIRECTION

SEVDIYOR IMOMKULOV, YULDASH SAIDOV

ABSTRACT. Let a function $u(x, y)$ be harmonic in the domain

$$D \times V_r = D \times \{y \in \mathbb{R}^m : |y| < r\} \subset \mathbb{R}^n \times \mathbb{R}^m$$

and for each fixed point x^0 from some a set $E \subset D$, the function $u(x^0, y)$, as a function of variable y , can be extended to a harmonic function on the whole $\mathbb{R}^m$. Then $u(x, y)$ harmonically extends to the domain $D \times \mathbb{R}^m$ as a function of variables x and y .

1. INTRODUCTION

The well known Hartogs lemma concerns the extension of holomorphic functions along fixed direction (see [1]) and states that if a function $f(z, w)$ is holomorphic in the domain

$$D \times \{w \in \mathbb{C} : |w| < r\} \subset \mathbb{C}^n \times \mathbb{C}$$

and for each fixed $z^0 \in D$ the function $f(z^0, w)$ of variable w holomorphically extends to the disk $\{w \in \mathbb{C} : |w| < R\}$, $R > r$, then $f(z, w)$ can be holomorphically extended to the domain $D \times \{w \in \mathbb{C} : |w| < R\}$ as a function of variables z and w .

Note that the Hartogs lemma remains to hold also for pluriharmonic and harmonic functions (see [3]) and can be formulated as follows. Let a function $u(x, y)$ be harmonic in the domain

$$D \times V_r = D \times \{y \in \mathbb{R}^m : |y| < r\} \subset \mathbb{R}^n \times \mathbb{R}^m,$$

and for each fixed $x^0 \in D$ the function $u(x^0, y)$ of variable y harmonically extends to the ball $V_R = \{y \in \mathbb{R}^m : |y| < R\}$, $R > r$. Then $u(x, y)$ harmonically extends to $D \times V_R$ as a function of variables x and y .

One of the main methods used in the investigation of harmonic extensions is to convert this problem to holomorphic extensions. For this purpose, the following lemma is proved in [3].

Lemma 1.1. *Consider the space $\mathbb{R}^n(x)$ embedded into $\mathbb{C}^n(z) = \mathbb{R}^n(x) + i\mathbb{R}^n(y)$, where $z = (z_1, z_2, \dots, z_n)$, $z_j = x_j + iy_j$, $j = 1, 2, \dots, n$. Then for each domain $D \subset \mathbb{R}^n(x)$ there exists some holomorphy domain $\hat{D} \subset \mathbb{C}^n(z)$, such that $D \subset \hat{D}$ and each harmonic function in D can be holomorphically extended to the domain $\hat{D}$, i.e., there exists a holomorphic function $f_u(z) \in \hat{D}$ such that $f_u|_D = u$.*

The main result of our paper reads as follows. The concept of N -sets of $Lh_0(D)$ which appears there is defined in the next section.

Theorem 1.2. *Let a function $u(x, y)$ be harmonic in the domain*

$$D \times V_r = D \times \{y \in \mathbb{R}^m : |y| < r\} \subset \mathbb{R}^n \times \mathbb{R}^m$$

1991 *Mathematics Subject Classification.* 32A65, 46J10, 46J15.

Key words and phrases. Harmonic function; harmonic extension; Hartogs theorem; Hartog's series; polar sets; holomorphic function; function class $Lh_0(D)$.

and for each fixed point x^0 from some set $E \subset D$, which is not embedded into a countable union of N -sets of $Lh_0(D)$, the function $u(x^0, y)$ of variable y can be extended to a harmonic function on the whole $\mathbb{R}^m$. Then $u(x, y)$ harmonically extends to the domain $D \times \mathbb{R}^m$ as a function of variables x and y .

2. The class of functions $Lh_0(D)$

Let $D \subset \mathbb{R}^n$ and $h(D)$ be the space of harmonic functions in D . By $Lh_\varepsilon(D)$ we denote the minimal class of functions which contains functions of the form $\alpha \ln |u(x)|$, $u(x) \in h(D)$, $0 < \alpha < \varepsilon$, and it is closed with respect to the operation of “upper regularization”, i.e., for any set of functions $u_\lambda(x) \in Lh_\varepsilon(D)$, $\lambda \in \Lambda$, where Λ is an index set, the function

$$\overline{\lim}_{y \rightarrow x} (\sup \{u_\lambda(y) : \lambda \in \Lambda\})$$

also belongs to the class $Lh_\varepsilon(D)$ ([4]).

The union of function classes $Lh_0(D) = \bigcup_{\varepsilon > 0} Lh_\varepsilon(D)$ we call *the set of Lh_0 -functions*.

Now, we will define N -set of the class $Lh_0(D)$. Let $\vartheta_k(x) \in Lh_0(D)$ be a monotonically increasing sequence of locally uniformly from above bounded functions. We denote

$$\vartheta(x) = \overline{\lim}_{y \rightarrow x} \lim_{k \rightarrow \infty} \vartheta_k(y) = x \in D.$$

Then everywhere in D holds the inequality

$$\lim_{k \rightarrow \infty} \vartheta_k(y) \leq \vartheta(x).$$

Definition 2.1. *The subsets of the set*

$$\{x \in D : \lim_{k \rightarrow \infty} \vartheta_k(y) < \vartheta(x)\}$$

are called N -sets of the class $Lh_0(D)$.

Note that if $\vartheta_k(x) \in Lh_0(D)$ is a sequence of locally uniformly from above bounded functions and

$$\overline{\lim}_{y \rightarrow x} \overline{\lim}_{k \rightarrow \infty} \vartheta_k(y) = \vartheta(x), \quad x \in D,$$

then the set

$$E = \{x \in D : \overline{\lim}_{k \rightarrow \infty} \vartheta_k(y) < \vartheta(x)\}$$

consists of countable union of N -sets of the class $Lh_0(D)$. Indeed, consider the sequence of functions

$$w_{l,j}(x) = \max_{l \leq k \leq j} \vartheta_k(x).$$

Clearly

$$\overline{\lim}_{k \rightarrow \infty} \vartheta_k(y) = \lim_{l \rightarrow \infty} \lim_{j \rightarrow \infty} w_{l,j}(x).$$

Since the sequence is increasing in j , we have

$$\lim_{j \rightarrow \infty} w_{l,j}(x) \leq \overline{\lim}_{l \rightarrow \infty} \lim_{j \rightarrow \infty} w_{l,j}(y), \quad x \in D,$$

and the sets

$$E_l = \{x \in D : \lim_{j \rightarrow \infty} w_{l,j}(x) < \overline{\lim}_{l \rightarrow \infty} \lim_{j \rightarrow \infty} w_{l,j}(y)\}, \quad l = 1, 2, \dots,$$

are N -set of class the $Lh_0(D)$. On the other hand, the sequences $\lim_{j \rightarrow \infty} w_{l,j}(x)$ and $\overline{\lim}_{l \rightarrow \infty} \lim_{j \rightarrow \infty} w_{l,j}(y)$, $j = 1, 2, \dots$, are monotonically decreasing, and

$$\overline{\lim}_{k \rightarrow \infty} \vartheta_k(x) = \lim_{l \rightarrow \infty} \lim_{j \rightarrow \infty} w_{l,j}(x) = \vartheta(x) = \lim_{l \rightarrow \infty} \overline{\lim}_{y \rightarrow x} \lim_{j \rightarrow \infty} w_{l,j}(y), \quad x \in D \setminus \bigcup_{l=1}^{\infty} E_l.$$

It follows that $E \subset \bigcup_{l=1}^{\infty} E_l$, i.e. $E = \bigcup_{l=1}^{\infty} (E_l \cap E)$.

Definition 2.2. *The set $E \subset D$ is called Lh_0 -polar with respect to D , if there exists a function $\vartheta(x) \in Lh_0(D)$ such that $\vartheta(x) \not\equiv -\infty$ and $\vartheta(x)|_E = -\infty$.*

Note that if $u(x) \in h(D)$, $u(x) \not\equiv 0$ and $E \subset \{u(x) = 0\}$, then E is Lh_0 -polar with respect to D .

Proposition 2.3. *Each Lh_0 -polar set with respect to D is contained in a countable union of N -sets of the class $Lh_0(D)$.*

Indeed, let E be a Lh_0 -polar set with respect to D . Then by Definition 2.2 there exists a function $\vartheta(x) \in Lh_0(D)$ such that $\vartheta(x) \not\equiv -\infty$ and $\vartheta(x)|_E = -\infty$. Consider the sequence of functions $\vartheta_k(x) = \frac{1}{k}\vartheta(x)$. Obviously, $\vartheta_k(x) \in Lh_0(D)$ and $\vartheta_k(x) \not\equiv -\infty$ and $\vartheta_k(x)|_E = -\infty$. Moreover, $\lim_{k \rightarrow \infty} \vartheta_k(x) = 0$ for almost all $x \in D$ and $\lim_{k \rightarrow \infty} \vartheta_k(x) = -\infty$ for all $x \in E$. It follows that

$$E \subset \left\{ x \in D : \lim_{k \rightarrow \infty} \vartheta_k(x) < \overline{\lim}_{y \rightarrow x} \lim_{k \rightarrow \infty} \vartheta_k(x) \right\}.$$

On the other hand, as we have showed above, the set

$$\left\{ x \in D : \lim_{k \rightarrow \infty} \vartheta_k(x) < \overline{\lim}_{y \rightarrow x} \lim_{k \rightarrow \infty} \vartheta_k(x) \right\}$$

consists of a countable union of N -sets of the class $Lh_0(D)$.

3. Proof of the theorem

Suppose that the number r is sufficiently large such that $\widehat{V} \subset \mathbb{C}^m$ contains the closure of the unit polydisc

$$U = \left\{ w = (w_1, w_2, \dots, w_m) \in \mathbb{C}^m : |w_1| < 1, |w_2| < 1, \dots, |w_m| < 1 \right\},$$

since the general case easily may be reduced to this case by linear changing of y . By the lemma of [3], for each fixed $x \in D$ the function $u(x, y)$ of variable y can be extended holomorphly to $\widehat{V}_r \subset \mathbb{C}^m$ and can be expanded into a Hartogs' series

$$u(x, w) = \sum_{|k|=0}^{\infty} c_k(x) w^k,$$

where $k = (k_1, k_2, \dots, k_m)$ is a multiindex. Clearly, the function $u(x, y)$ is harmonic in each variable and coefficients $c_k \in h(D)$. Also, according to the Cauchy's inequalities (see [1], [2]), for each set $D_0 : E \subset D_0 \subset\subset D$ the following estimation holds

$$|c_k(x)| \leq M, \quad \forall k, \quad \forall x \in D_0,$$

where $M = \sup\{|u(x, w)| : (x, w) \in \overline{D_0} \times \overline{U}\}$, i.e., the sequence of functions $c_k(x)$ is locally uniformly bounded in the domain D . On the other hand, by the same lemma

from [3], for each fixed $x \in E$ the function $u(x, w)$ of the variable w is holomorphic everywhere in $\mathbb{C}^m$, which means that for all $x \in E$

$$\overline{\lim}_{|k| \rightarrow \infty} \sqrt[k]{|c_k(x)|} = 0.$$

It follows that the sequence $\frac{1}{|k|} \ln |c_k(x)|$ is locally uniformly bounded from above and

$$\overline{\lim}_{|k| \rightarrow \infty} \frac{1}{|k|} \ln |c_k(x)| = -\infty, \quad \forall x \in E.$$

Put $\overline{\lim}_{y \rightarrow x} \overline{\lim}_{|k| \rightarrow \infty} \frac{1}{|k|} \ln |c_k(y)| = \vartheta(x)$. Since the set

$$F = \left\{ x \in D : \overline{\lim}_{|k| \rightarrow \infty} \frac{1}{|k|} \ln |c_k(x)| < \vartheta(x) \right\}$$

consists of the countable union of the N -sets of $Lh_0(D)$ and E is not contained in a countable union of the N -sets of $Lh_0(D)$, the set $E \setminus F = \{x \in E : \vartheta(x) = -\infty\}$ is not contained in a countable association of the N -sets of $Lh_0(D)$ as well. By Proposition 2.3 the set $E \setminus F$ is not Lh_0 -polar with respect to D . Consequently, $\vartheta(x) \equiv -\infty$, i.e.,

$$\overline{\lim}_{|k| \rightarrow \infty} \frac{1}{|k|} \ln |c_k(x)| = -\infty, \quad \forall x \in D.$$

Thus, we get

$$\overline{\lim}_{|k| \rightarrow \infty} \sqrt[k]{|c_k(x)|} = 0, \quad \forall x \in D.$$

Therefore, for all $x \in D$, the function $u(x, y)$ of variable y extends to a harmonic function on the whole $\mathbb{R}^m$. Hence, $u(x, y)$ harmonically extends in $D \times \mathbb{R}^m$. The theorem is proved.

REFERENCES

- [1] *B.W. Shabat* : Introduction to complex analysis. Part II. Nauka, Moskva, 1985. (In Russian)
- [2] *V.S. Vladimirov*: Methods of theory functions several complex variables. Moskva, Nauka 1964, p. 411.(In Russian)
- [3] *A.S. Sadullaev, S.A. Imomkulov*: Extension of holomorphic and pluriharmonic functions with thin singularities on parallel sections. Proceedings of the Steklov Institute of Mathematics, Moskva, (253), 2006, pp.158-174.
- [4] *V.P. Zahariuta*: Space of harmonic functions. In: Functional analysis, Lecture Notes in Pure and Applied Math., Dekker New York-Basel-Hong Kong, 1993, pp.497-522.

Authors' addresses: *S.A. Imomkulov, Y.R. Saidov*, Urganch State University, Urganch city, Khamid Alimjan -14, 740000, Usbekistan.

e-mail: (S.A. Imomkulov), sevdi@rambler.ru, (Y.R. Saidov) ysaidov@rambler.ru